arXiv:1109.2106v1 [math.GR] 9 Sep 2011

AUTOMORPHISM CLASSES OF ELEMENTS IN FINITELY GENERATED ABELIAN GROUPS.

CHARLES F. ROCCA JR.

Abstract. We will show that every element of a finitely generated abelian group is automorphically equivalent what we will define to be a *representative element* in a *repeat-free subgroup*, and for finite abelian groups we can count the number of automorphism classes of elements.

1. Representative Elements

A finite abelian p-group G_p can be written

$$G_p = \mathbb{Z}_{p^{r_1}}^{k_1} \oplus \cdots \oplus \mathbb{Z}_{p^{r_n}}^{k_n},\ r_i < r_{i+1}\ for\ all\ i.$$

However for this paper we will write such a group as follows

$$G_p = [\mathbb{Z}_{p^{r_1}} \oplus \cdots \oplus \mathbb{Z}_{p^{r_n}}] \oplus [\mathbb{Z}_{p^{r_1}}^{k_1-1} \oplus \cdots \oplus \mathbb{Z}_{p^{r_n}}^{k_n-1}],\ r_i < r_{i+1}.$$

This splits the group into two subgroups, a *repeat-free subgroup*,

$$G_p^{rf} = \mathbb{Z}_{p^{r_1}} \oplus \cdots \oplus \mathbb{Z}_{p^{r_n}},$$

which contains one copy of each of the factors $\mathbb{Z}_{p^{r_i}}$ and a *remainder subgroup*,

$$G_p^{rm} = \mathbb{Z}_{p^{r_i}}^{k_1-1} \oplus \cdots \oplus \mathbb{Z}_{p^{r_n}}^{k_n-1},$$

that contains the remaining factors of the group.

For the remainder of this article let G be a finitely generated abelian group of the form

$$G = G_{p_1} \oplus \cdots \oplus G_{p_m} \oplus \mathbb{Z}^l,$$

where l and $m \geq 0$ and the p_i are distinct primes with $p_i < p_{i+1}$ for all i. The above definition of a repeat free subgroup can be extended to any finitely generated abelian group G as follows.

Definition 1. *Given a finitely generated abelian group G a* repeat-free subgroup *has one of the following forms*

(1) $G^{rf} = G_{p_1}^{rf} \oplus \cdots \oplus G_{p_m}^{rf} \oplus \mathbb{Z}$ *if l and $m \geq 1$, or*
(2) $G^{rf} = G_{p_1}^{rf} \oplus \cdots \oplus G_{p_m}^{rf}$ *if $l = 0$ and $m \geq 1$, or*
(3) $G^{rf} = \mathbb{Z}$ *if $l \geq 1$ and $m = 0$.*

The next definition defines our set of *representative elements* in a repeat-free finite abelian p-group.

Definition 2. *An element $g = (g_1, \ldots, g_n)$ of a repeat-free finite abelian p-group $G_p^{rf} = \mathbb{Z}_{p^{r_1}} \oplus \cdots \oplus \mathbb{Z}_{p^{r_n}}$ $r_i < r_{i+1}$ for all i is a* representative element *if*

(1) *for all i g_i is either 0 or a power of p,*

2000 *Mathematics Subject Classification.* Primary 20K30, 20K01.

(2) *for all i, j if $i < j$, then $g_i < g_j$, and*
(3) *for all i, j if $i < j$, then the order of g_i in $\mathbb{Z}_{p^{r_i}}$ is less than the order of g_j in $\mathbb{Z}_{p^{r_j}}$.*

We will show that in a finite abelian p-group every element is automorphically equivalent to a representative element of a repeat-free subgroup. Therefore we let a *representative element* of a finite abelian p-group be an element of a repeat-free subgroup which is a representative element in that subgroup. Extending this definition to all finitely generated abelian groups we get.

Definition 3. *Let G ba a finitely generated abelian group, then we say $g = (g_1, g_2, \ldots, g_m, Z) \in G$ is a* representative element *if*

(1) *for all j, g_j is a representative element of G_{p_j},*
(2) *the element $Z \in \mathbb{Z}^l$ is of the form $(z, 0, \ldots, 0)$ and*
(3) *for all j, each term of g_j is relatively prime to z.*

So, the primary goal of this paper is the following theorem.

Theorem 1. *Every element of a finitely generated abelian group is automorphically equivalent to an unique representative element.*

The proof of this result has two main steps:

(1) Every element is automorphically equivalent to an element in a repeat-free subgroup.
(2) Every element of a repeat-free subgroup is automorphically equivalent to a representative element.

The following lemma, in which we think to endomorphisms of finite abelian groups as matrices, is a major tool in proving these results.

Lemma 1. [1] *An endomorphism φ of $G_p = \mathbb{Z}_{p^{r_1}} \oplus \mathbb{Z}_{p^{r_2}} \oplus \cdots \oplus \mathbb{Z}_{p^{r_m}}$, p prime and $r_i \leq r_{i+1}$ for all i, is an automorphism if and only if the reduction of φ modulo p, written φ_p, is an automorphism of $\mathbb{Z}_p^m$.*

The following proposition establishes the first step in the proof of the theorem.

Proposition 1. *Every element in a finitely generated abelian group G is automorphically equivalent to an element in the repeat-free subgroup. In particular:*

a) *Each element $g = (g_1, \ldots, g_n) \in \mathbb{Z}_{p^r}^n$, $n \geq 1$ is automorphically equivalent to an element of the form $(p^l, 0, \cdots, 0)$, where the order of g is p^{r-l}.*
b) *Each element $g = (g_1, \ldots, g_k)$ of $\mathbb{Z}^k$ is automorphically equivalent to one of the form $(d, 0, \cdots, 0)$, where $d = \gcd(g_1, \ldots, g_k)$.*

Proof: Let $g \in \mathbb{Z}_{p^r}^n$; each term in g is of the form $g_i = a_i p^{l_i}$, where a_i is relatively prime to p. Therefore g is mapped automorphically to the element $g' = (p^{l_1}, \cdots, p^{l_n})$ by multiplying each g_i by the inverse of a_i modulo p^r.

The order of g' which is the same as the order of g, is p^{r-l}. Therefore at least one term of g' is equal to p^l, without loss of generality let $g'_1 = p^l$. Also, every term of g' will be divisible by p^l, that is $l_i \geq l$ for all i. Therefore

$$\begin{bmatrix} 1 & 0 & \cdots & 0 \\ -p^{l_2-l} & 1 & \cdots & 0 \\ \vdots & & \ddots & \vdots \\ -p^{l_n-l} & 0 & \cdots & 1 \end{bmatrix} \begin{bmatrix} p^l \\ p^{l_2} \\ \vdots \\ p^{l_n} \end{bmatrix} = \begin{bmatrix} p^l \\ 0 \\ \vdots \\ 0 \end{bmatrix},$$

where it is clear from the lemma 1 that this matrix is an automorphism, thus proving part (a).
The element $g = (g_1, \ldots, g_k)$ of $\mathbb{Z}^k$ can be reduced using the Euclidian Algorithm by the repeated application of elementary row operations and this establishes (b). $\square$

To show that every element of a repeat-free subgroup is automorphically equivalent to a representative element we first restrict ourselves to a finite repeat-free p-group and define a *basic reduction* of an element.

Definition 4. *Given a repeat-free p-group*

$$G_p^{rf} = \mathbb{Z}_{p^{r_1}} \oplus \cdots \oplus \mathbb{Z}_{p^{r_n}}, \ r_i < r_{i+1} \ for\ all\ i$$

and an element $g = (p^{l_1}, \ldots, p^{l_n})$*, a* Basic Reduction about position i of g *is the following:*

for every $j \neq i$ *with* $p^{l_j} \geq p^{l_i}$ *and* $p^{r_j - l_j} \leq p^{r_i - l_i}$ *replace* p^{l_j} *with 0,*
i.e. if the j^{th} *term has a greater value but lesser order than the* i^{th} *term replace it by 0.*

If the number of non-zero terms decreases, then the basic reduction is a non-trivial basic reduction*, otherwise it is a* trivial reduction*. We will say that an element is* reduced *if there are no non-trivial basic reductions.*

The matrix for the reduction transformation has elements

$$a_{lk} = \begin{cases} 1 & if\ l = k \\ -p^{l_j - l_i} & if\ l = j\ and\ k = i \\ 0 & otherwise \end{cases}$$

which is similar to the matrix in Proposition 1 and is again by by Lemma 1 is an automorphism.

Lemma 2. *Let*

$$G_p^{rf} = \mathbb{Z}_{p^{r_1}} \oplus \cdots \oplus \mathbb{Z}_{p^{r_n}}, \ r_i < r_{i+1} \ for\ all\ i$$

be a repeat-free p-group. Then an element $g \in G_p^{rf}$ *is reduced if and only if g is a representative element.*

Proof: Let $g = (p^{l_1}, \ldots, p^{l_n})$ be a reduced element of G_p^{rf} and let $1 \leq i < j \leq n$.

Case 1: If $p^{l_i} \geq p^{l_j}$, then the order of p^{l_i} in $\mathbb{Z}_{p^{r_i}}$ is strictly less than the order of p^{l_j} in $\mathbb{Z}_{p^{r_j}}$. Therefore, we can perform a basic reduction about position j, contradicting the assumption that g is reduced.

Case 2: If $p^{l_i} < p^{l_j}$ and the order of p^{l_i} in $\mathbb{Z}_{p^{r_i}}$ is greater than or equal to the order of p^{l_j} in $\mathbb{Z}_{p^{r_j}}$, then we can perform a basic reduction about position i again contradicting the assumption that g was reduced. Therefore, if g in G_p^{rf} is reduced, then it is a representative element.

Now suppose that g is a representative element. Let $1 \leq i < j \leq n$ so that $p^{l_i} < p^{l_j}$ and the order of p^{l_i} in $\mathbb{Z}_{p^{r_i}}$ is strictly less than the order of p^{l_j} in $\mathbb{Z}_{p^{r_j}}$, when p^{l_i} and p^{l_j} are non-zero. If we performed a basic reduction about position i, then p^{l_j} would be unchanged since the order of p^{l_i} is less than that of p^{l_j}. Similarly, if we performed a basic reduction about position j, p^{l_i} would remain unchanged since $p^{l_i} < p^{l_j}$. Since i and j are arbitrary if g is a representative element, then g is reduced. $\square$

We are now in a position to prove the main result for the case of repeat-free p-groups.

Proof: (Theorem 1 for repeat-free p-groups) Let $g = (g_1, \ldots, g_n)$ be an element in a finite repeat-free p-group G_p^{rf}. Applying Proposition 1 we know that g is automorphically equivalent to some g' in which each term is a power of p. Now, if g' is not a reduced element, then we can perform a non-trivial basic reduction and the the resulting element, which is automorphically equivalent to g', will have strictly fewer non-zero terms. Since the rank of G_p^{rf} is finite the process of making non-trivial reductions will terminate, and the resulting element will be a reduced element and therefore a representative element.

Finally we show that representative elements of automorphism classes are unique.

Claim 1. *If g and h are automorphically equivalent representative elements in G_p, then $g = h$.*

Proof: (of claim) Let $g = (g_1, \ldots, g_n)$, $h = (h_1, \ldots, h_n)$ and let A be the matrix representing the automorphism taking h to g. Suppose $h_i \neq 0$ and let p^{l_i}, $0 \leq l_i \leq r_i$ be the order of h_i in $\mathbb{Z}_{p^{r_i}}$ so that we may write $h_i = p^{r_i - l_i}$. By Lemma 1 the automorphism A must have the form

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ p^{r_2 - r_1} a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & & & \vdots \\ p^{r_n - r_1} a_{n1} & p^{r_n - r_2} a_{n2} & \cdots & a_{nn} \end{bmatrix}$$

where a_{ii} is relatively prime to p. Therefore, assuming that $h_i \neq 0$,

$$g_i = p^{r_i - l_1} a_{i1} + \cdots + p^{r_i - l_i} a_{ii} + \cdots + p^{r_n - l_n} a_{in}.$$

Since h is a representative element, for any non-zero term h_j if $j < i$, then the order of h_j in $\mathbb{Z}_{p^{r_j}}$ is less than the order of h_i in $\mathbb{Z}_{p^{r_i}}$; i.e. $l_j < l_i$. Hence, $r_i - l_j > r_i - l_i$ and $p^{r_i - l_i}$ divides $p^{r_i - l_j}$. If $j > i$, then $h_i < h_j$ and $p^{r_i - l_i}$ divides $p^{r_j - l_j}$. Thus

$$g_i = p^{r_i - l_i}(p^{l_i - l_1} a_{i1} + \cdots + a_{ii} + \cdots + p^{r_n - l_n - (r_i - l_i)} a_{ni}).$$

Since a_{ii} is relatively prime to p

$$b_i = (p^{l_i - l_1} a_{i1} + \cdots + a_{ii} + \cdots + p^{r_n - l_n - (r_i - l_i)} a_{ni}).$$

is also relatively prime to p. Thus, when h_i is non-zero, $g_i = b_i h_i$, b_i relatively prime to p. However g is a representative element therefore g_i is a power of p, $b_i = 1$ and $g_i = h_i$.

An identical argument shows that $h_i = g_i$ when g_i is non-zero. Therefore for all i, $g_i = h_i$ and so $g = h$. □

If G_p is any finite abelian p-group, then every element of G_p is automorphically equivalent to a unique representative element in G_p^{rf}, which is also a representative element for G_p. Finally, if G is any finitely generated abelian group, then we have established that any element of G is automorphically equivalent to element satisfying the first two conditions of the definition of representative element. It can be shown that any element satisfying the first two conditions is automorphically equivalent to one that satisfies the third condition and has the same number or fewer nonzero terms. As an illustration consider the following example.

Let $G^{rf} = \mathbb{Z}_{p^{r_1}} \oplus \mathbb{Z}_{p^{r_2}} \oplus \mathbb{Z}$ and let $g = [p^{l_1}, p^{l_2}, kp]$ be any element of G^{rf} satisfying the first two conditions of the definition of a representative element but not the third. Then

$$\begin{bmatrix} 1 & 0 & p^{r_1-1}-p^{l_1-1} \\ 0 & 1 & p^{r_2-1}-p^{l_2-1} \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} p^{l_1} \\ p^{l_2} \\ kp \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ kp \end{bmatrix}$$

and so g is automorphically equivalent to an element satisfying the third condition of our definition and has strictly fewer nonzero terms.

2. Counting Automorphism Classes

Having completed the main result we will now show how to count the number of representative elements in a given finite abelian group.

Let $G_p^{rf} = \mathbb{Z}_{p^{r_1}} \oplus \mathbb{Z}_{p^{r_2}} \oplus \cdots \oplus \mathbb{Z}_{p^{r_k}}$ where $r_i < r_{i+1}$ for all i be a repeat-free p-group and $g = (p^{l_1}, p^{l_2}, \ldots, p^{l_k})$ be an automorphism class representative in G_p^{rf}. The non-zero terms of g are both increasing and order increasing, thus for $0 \leq i < j \leq k$ if p^{l_i} and p^{l_j} are non-zero terms of g we know that

$$p^{l_i} < p^{l_j}$$

and

$$p^{r_i - l_i} < p^{r_j - l_j}.$$

Hence,

$$l_i < l_j < l_i + (r_j - r_i).$$

We shall refer to the value $r_j - r_i$ as the *gap* between the terms $\mathbb{Z}_{p^{r_i}}$ and $\mathbb{Z}_{p^{r_j}}$ of G_p^{rf}, and in particular we will denote $r_{i+1} - r_i$ by n_i. As an immediate consequence of the above conclusion we know that if p^{l_i} is non-zero, then either p^{l_j} is zero or we have at most $r_j - r_i - 1$ choices for l_j. Further, if $j = i+1$ and $n_i = r_{i+1} - r_i = 1$, then one or the other of these two terms must be zero. Therefore, if the gap between two terms of a group is one, then in every automorphism class representative at least one of the terms will be zero.

For motivation let us count the number of automorphism class representatives in $G_p^{rf} = \mathbb{Z}_{p^{r_1}} \oplus \mathbb{Z}_{p^{r_2}} \oplus \mathbb{Z}_{p^{r_3}}$ where $r_1 < r_2 < r_3$. We begin by counting the number of automorphism classes with a given number of non-zero terms.

# of Non-Zero Terms	# of Automorphism Classes
0	1
1	$r_1 + r_2 + r_3$
2	$r_1(n_1 - 1) + r_1(r_3 - r_1 - 1) + r_2(n_2 - 1)$
3	$r_1(n_1 - 1)(n_2 - 1)$

However $r_2 = r_1 + n_1$ and $r_3 = r_1 + n_1 + n_2$ therefore the above equations can be rewritten as,

# of Non-Zero Terms	# of Automorphism Classes
0	1
1	$r_1 + (r_1 + n_1) + (r_1 + n_1 + n_2)$
2	$r_1(n_1 - 1) + r_1(n_1 + n_2 - 1) + (r_1 + n_1)(n_2 - 1)$
3	$r_1(n_1 - 1)(n_2 - 1)$

which gives us,

Non-Zero Terms	# of Automorphism Classes
0	1
1	$3r_1 + 2n_1 + n_2$
2	$r_1(2n_1 + 2n_2 - 3) + n_1 n_2 - n_1$
3	$r_1(n_1 n_2 - n_1 - n_2 + 1)$

and the number of automorphism classes of elements equals

$$r_1(1 + n_1 + n_2 + n_1 n_2) + 1 + n_1 + n_2 + n_1 n_2 = (r_1 + 1)(n_1 + 1)(n_2 + 1).$$

Proposition 2. *The number of automorphism classes of elements in a repeat-free finite abelian p-group*

$$G_p^{rf} = \mathbb{Z}_{p^{r_1}} \oplus \mathbb{Z}_{p^{r_2}} \oplus \cdots \oplus \mathbb{Z}_{p^{r_k}},\ r_i < r_{i+1}\ \mathit{for\ all}\ i$$

is equal to

$$(r_1 + 1)(n_1 + 1)(n_2 + 1) \cdots (n_{k-1} + 1),$$

where

$$n_i = r_{i+1} - r_i.$$

Proof: We have shown above that this is true for a group with three terms. We proceed by induction on the number of terms in G_p^{rf}. By induction the group

$$H_p^{rf} = \mathbb{Z}_{p^{r_1}} \oplus \mathbb{Z}_{p^{r_2}} \oplus \cdots \oplus \mathbb{Z}_{p^{r_{k-1}}}$$

has $(r_1 + 1)(n_1 + 1)(n_2 + 1) \cdots (n_{k-2} + 1)$ automorphism classes of elements. This is also the number of automorphism class representatives in G_p^{rf} in which the k^{th} term is 0. Therefore we complete the proposition with the following claim.

Claim 2. *If $1 < j \leq k$, then in*

$$G_p^{rf} = \mathbb{Z}_{p^{r_1}} \oplus \mathbb{Z}_{p^{r_2}} \oplus \cdots \oplus \mathbb{Z}_{p^{r_k}}$$

the number of automorphism class representatives in which the j^{th} term is the last non-zero term is

$$(r_1 + 1)(n_1 + 1)(n_2 + 1) \cdots (n_{j-2} + 1)n_{j-1}.$$

Proof: (of claim) If $j = 2$, then there are r_2 automorphism class representatives in which the second term is the only non-zero term and $r_1(n_1 - 1)$ in which the first and second terms are the only non-zero terms. Therefore the number of automorphism class representatives in which the second term is the last non-zero term is

$$\begin{aligned} r_1(n_1 - 1) + r_2 &= r_1 n_1 - r_1 + r_1 + n_1 \\ &= (r_1 + 1)n_1. \end{aligned}$$

Now let $2 < j \leq k$, if $2 \leq i < j$ and the i^{th} term is the last non-zero term of an element prior to the j^{th} term, then there are $(r_j - r_i - 1)$ non-zero choices for the value of the j^{th} term and by induction

$$(r_1 + 1)(n_1 + 1)(n_2 + 1) \cdots (n_{i-2} + 1)n_{i-1}(r_j - r_i - 1)$$

automorphism class representatives in which the j^{th} term is the last non-zero term and the i^{th} term is the second to last non-zero term. So letting

$$A_i = (r_1 + 1)(n_1 + 1)(n_2 + 1) \cdots (n_{i-2} + 1)n_{i-1}$$

and summing over all i between 2 and j, the total number of automorphism classes in which the j^{th} term is the last non-zero term is

$$A_j = r_j + r_1(r_j - r_1 - 1) + \sum_{i=2}^{j-1} A_i(r_j - r_i - 1).$$

However for all $1 \leq i < j \leq k$:

$$r_j = r_{j-1} + n_{i-1}$$

and

$$(r_j - r_i - 1) = (r_{j-1} - r_i - 1 + n_{j-1})$$

so we may rewrite the previous equation as

$$(r_{j-1} + n_{j-1}) + (r_1(r_{j-1} - r_1 - 1) + r_1 n_{j-1}) + \sum_{i=2}^{j-1}(A_i(r_{j-1} - r_i - 1) + A_i n_{j-1})$$

And since

$$A_{j-i} = r_{j-1} + r_1(r_{j-1} - r_1 - 1) + \sum_{i=2}^{j-2} A_i(r_{j-1} - r_i - 1)$$

we get

$$\begin{aligned}
A_j &= A_{j-1} + n_{j-1} + r_1 n_{j-1} + A_{j-1}(r_{j-1} - r_{j-1} - 1) + \sum_{i=2}^{j-1} A_i n_{j-1} \\
&= A_{j-1} - A_{j-1} + \left(r_1 n_{j-1} + n_{j-1} + \sum_{i=2}^{j-1} A_i n_{j-1} \right) \\
&= r_1 n_{j-1} + n_{j-1} + \sum_{i=2}^{j-1} A_i n_{j-1} \\
&= n_{j-1} \left((r_1 + 1) + \sum_{i=2}^{j-1} A_i \right) \\
&= (r_1 + 1)(n_1 + 1)(n_2 + 1) \cdots (n_{j-2} + 1) n_{j-1},
\end{aligned}$$

thus proving the claim. In order to finish the proposition we observe that, by induction, the number of automorphism classes in which the k^{th} term is zero is

$$(r_1 + 1)(n_1 + 1)(n_2 + 1) \cdots (n_{k-2} + 1)$$

and from the claim the number of automorphism classes in which the k^{th} term is non-zero is

$$(r_1 + 1)(n_1 + 1)(n_2 + 1) \cdots (n_{k-2} + 1) n_{k-1},$$

and the sum of these two gives the desired result. □

Since automorphisms respect the prime decomposition of finite abelian groups we get this final general result.

Theorem 2. *The number of automorphism classes of elements in a finite abelian group*

$$G = G_{p_1} \oplus \cdots \oplus G_{p_m}$$

where

$$G_{p_j} = \mathbb{Z}_{p_j^{r_1}}^{k_1} \oplus \cdots \oplus \mathbb{Z}_{p_j^{r_n}}^{k_n},\ r_i < r_{i+1}$$

is equal to the product of the number of automorphism classes in each individual $G_{p_j}^{rf}$.

REFERENCES

[1] C.F. Rocca Jr. and E.C. Turner, *Test Elements in Finitely Generated Abelian Groups*, Int. J. Algebra and Computation, Vol. 12, No. 4 (2002) pp.569-573.

[2] J.C. O'Neill and E. C. Turner, *Test elements in direct products*, Int. J. Algebra and Comput, Vol. 10, No. 6 (2000), pp.751-756.

Current address: Department of Mathematics, Western Connecticut State University, Danbury, CT 06810

E-mail address: `roccac@wcsu.edu`